

The Exotic Molecular Ion H_4^{3+} in a Strong Magnetic Field

H.Olivares Pilón

Instituto de Ciencias Nucleares, UNAM, Apartado Postal 70-543, 04510 México

Abstract

Using the variational method, a detailed study of the lowest $m = 0, -1$ electronic states of the exotic molecular ion H_4^{3+} in a strong magnetic field, in the linear symmetric configuration parallel to the direction of the magnetic field is carried out. A extended study of the $1\sigma_g$ ground state (J.C. López and A. Turbinder, *Phys. Rev. A* **62**, 022510, 2000) was performed obtaining that the potential energy curve displays a sufficiently deep minimum for finite internuclear distances, indicating the possible existence of the molecular ion H_4^{3+} , for magnetic fields of strength $B \gtrsim 3 \times 10^{13}$ G. It is demonstrated that the excited state $1\pi_u$ can exist for a magnetic field $B = 4.414 \times 10^{13}$ G corresponding to the limit of applicability of the non-relativistic theory.

Keywords: Strong Magnetic Field; Molecular Ion H_4^{3+} ; Excited States.

PACS numbers: 31.15.Pf, 31.10.+z, 97.10.Ld

I. INTRODUCTION

The neutron stars were predicted by L. Landau [1] in 1932. This kind of stars are characterized by an enormous magnetic field ($B \sim 10^{12}$ G -10^{13} G). Since the discovery of neutron stars in 1967 the behavior of the molecular and atomic systems in such strong magnetic field has been object of intense investigations. The first studies were made in plasma physics and astrophysics by Kadomstev-Kudrayavstev [2] and Ruderman [3], respectively. The results demonstrated that the binding energy of a molecular system grows substantially with magnetic field increase, while the size of the systems decreases. As a consequence of such a behavior, the formation of a molecular chain aligned in the magnetic field direction arise.

Recent studies of the Coulomb systems in a strong magnetic field have been encouraged by the observation of the neutron star 1E1207.4-5209 (*Chandra* X-ray observatory 2002). These observations displayed two absorption features at 0.7 keV and 1.4 keV. The origin of such lines could be associate to the presence of exotic molecular systems in the atmosphere of the above mentioned star [4]. In particular, the studies of one-electron molecular systems, traditional and exotic have been intensified, such as H_2^+ , H_3^{2+} , H_4^{3+} , $(HeH)^{2+}$, He_2^{3+} y Li_2^{5+} (see for example [5]).

The focus of this article is a quantitative analysis of the lowest electronic states of the molecular system H_4^{3+} placed in a strong magnetic field. The variational method in the framework of non-relativistic theory is used, *i.e.* $B \leq 4.414 \times 10^{13}$ G (Schwinger limit). Specifically, an extension of the preliminary study [6] for the ground state is made, and the possible existence of excited states is carried out.

Atomic units are used throughout ($\hbar = e = m_e = 1$) and the energy is expressed in Rydbergs (Ry). Spectroscopic convention is used to denote the states. A number indicates the principal quantum number, Greek let-

ter $\sigma, \pi, \delta \dots$ indicate the magnetic quantum number for $|m| = 0, 1, 2 \dots$ respectively. The spatial parity is defined by g (gerade) for states with even parity and u (ungerade) for odd parity states. Particularly, the ground state is indicated as $1\sigma_g$.

II. THE MOLECULAR ION H_4^{3+}

The exotic molecular ion H_4^{3+} is a system constituted by four protons and one electron. For the present work the protons are consider over a straight line (linear configuration) parallel to the lines of magnetic field B (parallel configuration). The magnetic field B is taken as constant and homogeneous in z direction. The Born-Oppenheimer approximation of zero order is assumed (protons are consider infinitely massive). The positions of the four protons over z axis are denoted as R_1 , R_2 , $-R_3$ and $-R_4$. We will suppose also that $R_1 = R_3$ and $R_2 = R_4$ (symmetric configuration (see Fig.1)).

In order to describe the magnetic field $\mathbf{B} = (0, 0, B)$, the vector potential in the symmetric gauge $\mathbf{A} = \frac{B}{2}(-y, x, 0)$ is assumed. The Hamiltonian that describes the molecular ion H_4^{3+} (in cylindrical coordinates) is

$$\hat{\mathcal{H}} = -\Delta + \hat{l}_z B + \frac{B^2 \rho^2}{4} + V(\rho, z), \quad (1)$$

where $\rho^2 = x^2 + y^2$ is the distance from the electron to the z axis. $\hat{l}_z$ is the z component of the angular momentum (which is a integral of motion with eigenvalue m -magnetic quantum number). $V(\rho, z)$ is the Coulomb potential of the interaction between the electron and each charged center, plus the classic repulsion term between




FIG. 1: The molecular ion H_4^{3+} in a linear configuration placed in a constant and homogeneous magnetic field directed according to z axis, $\mathbf{B} = (0, 0, B)$. The protons positions are indicated by the four points over z axis. The position of the electron is marked by a point out of the axis.

the charged centers. The potential is given by

$$V(\rho, z) = -\frac{2}{r_1} - \frac{2}{r_2} - \frac{2}{r_3} - \frac{2}{r_4} + \frac{2}{R_2 - R_1} + \frac{2}{R_2 + R_3} + \frac{2}{R_2 + R_4} + \frac{2}{R_1 + R_3} + \frac{2}{R_1 + R_4} + \frac{2}{R_4 - R_3}, \quad (2)$$

where $r_i = \sqrt{\rho^2 + (z - z_i)^2}$ is the distances between the electron and the corresponding proton (z_i are the coordinates of each proton over the z axis).

In the Hamiltonian (1) the spin contribution is neglected because it only changes a reference level for the total energy.

III. THE TRIAL FUNCTIONS

In order to study the system H_4^{3+} the variational method is used. For the choosing of the trial functions, the fitting criteria used by Turbner-López was followed (see for example [5]). These criteria have proved to give much precise results, qualitatively and quantitatively, for different Coulomb systems in strong magnetic fields. Basically these are:

- 1.- The potential $V_t = \frac{\Delta\psi_t}{\psi_t}$, for which the trial function ψ_t is an exact eigenfunction of $\hat{\mathcal{H}}_t = -\Delta + V_t$, with zero energy, should reproduce the original potential near the Coulomb singularities and the harmonic oscillator behavior at large distances.
- 2.- The symmetries of the system must be included.
- 3.- For the ground state, the trial function should not vanish inside the domain where the problem is defined (Theorem of Perron).

In addition, since the charged centers (protons) are identical, the trial function is chosen symmetric with respect to the interchange of these, *i.e.*, invariant under transformations $P_{ij}(r_i \leftrightarrow r_j)$, $i, j = 1 \dots 4$.

The trial function is a linear combination of terms that fulfill the requirements above mentioned. The general form of the *ansatz* is

$$\Psi_m \equiv \mathcal{U}_m \Psi_0, \quad (3)$$

where

$$\Psi_0 = \left(\sum_{perm\{\alpha_1, \dots, \alpha_4\}} e^{-\alpha_1 r_1 - \alpha_2 r_2 - \alpha_3 r_3 - \alpha_4 r_4} \right) e^{-\beta B \frac{\rho^2}{4}}, \quad (4a)$$

$$\mathcal{U}_m = e^{im\phi} \rho^{|m|}. \quad (4b)$$

Here Ψ_0 corresponds to the general trial function for the ground state ($m = 0$). $\alpha_{1,2,3,4}$ and β are variational parameters. The sum is over all permutations of the parameters $\alpha_{1,2,3,4}$. Excitations for different values of the magnetic quantum number m are included with the $\mathcal{U}_m$ term. The trial functions Ψ_m given by Eq. (3) corresponding to different values of m are orthogonal and result suitable to describe the lowest states inside each family of own states with quantum number m defined.

A. Gauge Rotation

Considering the trial function Ψ_m given by Eq. (3) and the Hamiltonian $\hat{\mathcal{H}}$ (see Eq. (1)), the total energy (for a state with a quantum magnetic number m) is

$$E_{T_{var}} = \min_{\{\alpha'_s, \beta'_s\}} \frac{\int \rho^{2|m|} \Psi_0 \hat{h}_m \Psi_0 d\mathbf{r}}{\int \rho^{2|m|} \Psi_0 \Psi_0 d\mathbf{r}}, \quad (5)$$

where

$$\hat{h}_m \equiv \mathcal{U}_m^{-1} \hat{\mathcal{H}} \mathcal{U}_m = \hat{\mathbf{p}}_m^2 + Bm + \frac{B^2}{4} \rho^2 + V(\rho, z), \quad (6)$$

is the gauge rotated Hamiltonian, and $\hat{\mathbf{p}}_m \equiv \mathcal{U}_m^{-1} \hat{\mathbf{p}} \mathcal{U}_m$ is the covariant moment. From here on, $E_{T_{var}}$ will denote simply as E_T . This formulation is very suitable since each Hamiltonian $\hat{h}_m$ (for m constant) describes a family of eigenstates with the quantum number m .

Moreover, the ground state and the excited states are described by trial functions which have the same structure (given by the function (4a)). The only difference appears as a weight factor $\rho^{2|m|}$.

The potential associated to an arbitrary trial function ϕ_0 is now determined by

$$V_0^{(m)} = \frac{\hat{\mathbf{p}}_m^2 \Phi_0}{\Phi_0} \equiv \frac{\tilde{\Delta} \Phi_0}{\Phi_0}, \quad (7)$$

where the Laplacian covariant $\tilde{\Delta}$ is defined as

$$\tilde{\Delta} = \Delta + 2|m| \left(\frac{1}{\rho} \frac{\partial}{\partial \rho} + \frac{i}{\rho^2} \frac{\partial}{\partial \phi} \right). \quad (8)$$

Hence, the general form of the potential associated with each trial function term Ψ_0 , (Eq. (4a)) is [11]

$$V_0^{(m)} = \left\{ \beta B \rho^2 - 2 \right\} \sum_{i=1}^4 \frac{\alpha_i}{r_i} + \sum_{i,k=1}^4 \alpha_i \alpha_k (\hat{\mathbf{n}}_i \cdot \hat{\mathbf{n}}_k) \quad (9)$$

$$- \beta B + \frac{(\beta)^2 B^2}{4} \rho^2 - 2|m| \sum_{i=1}^4 \frac{\alpha_i}{r_i} - |m| \beta B,$$

where

$$\hat{\mathbf{n}}_i \cdot \hat{\mathbf{n}}_k = \frac{1}{r_i r_k} \{ \rho^2 + (z - z_i)(z - z_k) \},$$

and $\hat{\mathbf{n}}_i \equiv (\mathbf{r} - \mathbf{r}_i)/|\mathbf{r} - \mathbf{r}_i|$ is a unitary vector oriented according to the line which connects the i th position charged center, with the position of electron $\mathbf{r}$. The potential (9) exactly reproduces the original potentials of Coulomb for $\alpha_1 = \alpha_2 = \alpha_3 = \alpha_4 = 1$, and of harmonic oscillator for $\beta = 1$.

IV. TRIAL FUNCTION TERMS Ψ_p

The election of the most general trial function Ψ_0 (Eq. (4a)) was in no way arbitrary. The α parameters may be interpreted as the effective charge of the protons and β parameter as the effective coupling of the electron with magnetic field B . With this characterization of the variational parameters, it is possible obtain different trial functions each one with certain physical interpretation.

Let us consider for example the case in which the interaction of the electron with each one of the protons is *coherent*. That is, the electron interacts in the same way with each proton. To describe such physical situation, it is enough to take $\alpha_1 = \alpha_2 = \alpha_3 = \alpha_4 \equiv \alpha_1$ in the Eq. (4a). Defining this function as $\psi_0^{(1)}$ the following is obtained

$$\psi_0^{(1)} = e^{-\alpha_1(r_1+r_2+r_3+r_4)} e^{-\beta_1 \frac{B}{4} \rho^2}.$$

This function is the Heitler-London type function (used to describe the system H_2^+ without magnetic field) multiplied by the lowest Landau orbital. The results of the calculus verify (a posteriori) that this function gives the major contribution in the description of the system near its equilibrium position.

In order to describe the physical situation at large internuclear distances, all α' s parameters are equal to zero, except one, which is denote as α_2 . The trial function is

$$\psi_0^{(2)} = \left\{ e^{-\alpha_2 r_1} + e^{-\alpha_2 r_2} + e^{-\alpha_2 r_3} + e^{-\alpha_2 r_4} \right\} e^{-\beta_2 \frac{B}{4} \rho^2},$$

which corresponds to a Hund-Mulliken type function (used to describe the system H_2^+ without magnetic field) multiplied by the lowest Landau orbital. This function describes the interaction of the electron with each of the protons independently (*incoherent interaction*).

It is now clear that different degenerations from the general function (Eq. (4a)) lead to different physical descriptions of the system, and therefore to several trial functions. In this case and in order not to discriminate between one and another of the different descriptions, the linear combination of all the possible particular cases was considered as trial function. Explicitly, the trial function used in the current study is

$$\Psi_t = A_1 \psi_0^{(1)} + A_2 \psi_0^{(2)} + A_3 \psi_0^{(3)} + A_4 \psi_0^{(4)} + \quad (10)$$

$$A_5 \psi_0^{(5)} + A_6 \psi_0^{(6)} + A_7 \psi_0^{(7)} + A_8 \psi_0^{(8)} +$$

$$A_9 \psi_0^{(9)} + A_{10} \psi_0^{(10)} + A_{11} \psi_0^{(11)}.$$

All the terms included in the last equation admit a certain physical interpretation. This are summarized in Table I where we also present the variational parameters number contained in each one of them. A detailed analysis of each term appears in [7].

Function	N.V.P.	Physical Description
$\psi_0^{(1)}$	2	Coherent Interaction
$\psi_0^{(2)}$	2	Incoherent Interaction
$\psi_0^{(3)}$	2	$H_2^+ + 2p$
$\psi_0^{(4)}$	2	$H_3^{2+} + p$
$\psi_0^{(5)}$	3	" $H_2^+ + H_2^+$ "
$\psi_0^{(6)}$	3	" $H_3^{2+} + H$ "
$\psi_0^{(7)}$	4	" $H_2^+ + H + H$ "
$\psi_0^{(8)}$	3	$p + "H_2^+ + H"$
$\psi_0^{(9)}$	4	$p + "H + H + H"$
$\psi_0^{(10)}$	3	$p + p + "H + H"$
$\psi_0^{(11)}$	5	" $H + H + H + H$ "

TABLE I: The trial function terms Ψ_t (Eq. 10) along with the number of variational parameters (N.V.P.) and its physical interpretation (see text).

Another degeneration of the Eq. (4a) happens when α 's parameters are equal by pairs, $\alpha_1 = \alpha_2 = \alpha$; $\alpha_3 = \alpha_4 = \alpha'$, ($\alpha \neq \alpha'$). The physical picture correspond to an electron shared by two protons of different form. The situation may be interpreted as an interaction $H_2^+ + H_2^+$ between two Hydrogen molecular ions, even though of course we only have one electron (these type of physical interpretations are denoted between quotation marks on Table I). In the same way, in the most general case (all α 's parameters are different), the corresponding function is a sum of products of four Hydrogen functions by one Landau function (see Eq. (4a)). This situation describes

a mixed state of four Hydrogen atoms sharing only one electron.

Along with the $A_{1,\dots,11}$ coefficients of the linear combination (Eq. (10)) and the positions $R_{1,\dots,4}$ (see Fig. 1) there is a total of 48 variational parameters. The normalization condition of the trial function allows to keep one of the A coefficients as constant. The symmetry restriction ($R_3 = R_1, R_4 = R_2$) fixes two parameters. Finally the number of effective variational parameters is reduced to 45.

V. COMPUTER TECHNIQUE

All calculations were made in a DELL PC with two Xeon processors of 2.8 GHz each. For minimization calculations the minimization routine MINUIT of the CERN-LIB library was used. Numerical integration is carried out using the routine D01FCF from the NAG-LIB library. The relative precision in the numerical integration was of 10^{-11} . It is important to mention that once all variational parameters are determined, the calculation of the variational energy takes a few seconds.

A. Minimization Strategy

The variational energy calculation includes two technical aspects: the numerical evaluation of integrals in two dimensions and minimization of energy with respect to the variational parameters. The objective in the minimization process is to find a point in the multidimensional parametric space that minimizes the total energy of the system (Eq. (5)). Since we have 45 variational parameters, the search of the minimum is a formidable, complicated and tedious task. For this reason it is not a good idea to start the procedure making a minimization in which all parameters vary simultaneously. The strategy used consisted in making minimization sequences in the following way

- Ansatz by Ansatz, knowing clearly that between the eleven terms that appear in the Eq. (10), the most general $\psi_{p0}^{(11)}$ should give us the best (lowest) value in total energy.
- Combinations of two by two Ansätze, taking ψ_{p0}^{11} as the common Ansatz to all these combinations.
- Some combinations of three by three Ansätze, taking as ground the previous combinations, where one more Ansatz was included.
- Beginning from that combination of three Ansätze with the best total energy value, new Ansätze were incorporated until there was a total of eleven of them.

Each step in a particular minimization consists in keeping some parameters as constant while others were varied. The election of which parameters must be kept constant and which free was always based on the physical interpretation of such parameters. In no way the calculation is automatic, rather, a sophisticated artisan treatment is required. With this procedure, a variational parameter configuration was obtained in which the systems total energy value is minimum. Each minimization procedure for a given magnetic field as a general statement took various months of calculations (see [7]).

VI. RESULTS

In this section the total energy E_T , the binding energy [12] E_b ($E_b \equiv B - E_T$), and the equilibrium distance of protons ($R_3 = R_1, R_4 = R_2$) for the ground state $1\sigma_g$ and the excited state $1\pi_u$ of the molecular ion H_4^{3+} are presented. The trial function used is shown in Eq. (10). For a comparison with the results obtained by Turbinder-López [13] the calculations were carried out considering two different conversion factors to atomic units. These factors are $B_0 = 2.35 \times 10^9$ G (the one adopted in this work) and $B_0 = 2.3505 \times 10^9$ G (the one adopted in [6]).

A. The Ground State $1\sigma_g$

For the ground state $1\sigma_g$ each Ansatz presents a minimum in the energy curve for the finite parameter values $R_{1,\dots,4}$. In Table II the results are present. The results carried out by Turbinder-López [6] (also present in Table II) only consider the four terms that appear in Table I.

A total energy curve of the system as function of R_2 was obtained (see Fig. 2). For this the parameter R_1 is considered constant to the corresponding value of the minimum total energy configuration (in symmetric configuration). The R_3 and R_4 parameters are now considered as variational parameters. In other words the symmetry restriction is eliminated. The curve presents a well defined minimum for finite internuclear distances in the symmetric configuration. The height of the barrier is $\Delta E \equiv E_{max} - E_{min} = 0.242$ Ry. The asymptotic behavior of this curve for large R_2 shows that $R_1 + R_3 \approx 0.1113$ (see Fig. 1) and $R_4 - R_3 \approx 0.1110$. These values are very similar to the equilibrium distances of the ion H_3^{2+} (see Table IV). This indicates that a decay $H_4^{3+} \rightarrow H_3^{2+} + p$ is possible. The energy as function of R_1, R_2, R_3 and R_4 , generates a surface. In particular the curve presented in Fig. 2 corresponds to the cut throughout the valley of this surface.

The electronic distribution corresponding to ground state $1\sigma_g$ is shown in Fig. 3. We can observe that the electronic cloud is mainly concentrated around the intermediate point to the protons.

$B(\text{G})$	$B_0(\times 10^9 \text{ G})$	$E_T(\text{Ry})$	$E_b(\text{Ry})$	$R_1(\text{a.u.})$	$R_2(\text{a.u.})$	
3×10^{13}	2.35	12727.5422	38.4153	0.0635	0.2763	Present
	2.35	18739.2526	43.7261	0.0574	0.2336	Present
4.414×10^{13}	2.3505	18735.2602	43.7230	0.0574	0.2336	Present
	2.3505	18735.4675	43.5157	0.0572	0.2326	Turbiner-López ^a

^aNot published.

TABLE II: Results obtained for ground state $1\sigma_g$ of the molecular ion H_4^{3+} in the presence of a magnetic field B for different values of the conversion factor B_0 and in comparison with those obtained by Turbiner-López.




FIG. 2: Total energy curve of the molecular ion H_4^{3+} in the ground state $1\sigma_g$ as a function of parameter R_2 , which presents a minimum for finite equilibrium distances. This in linear configuration and parallel to the magnetic field $B = 4.414 \times 10^{13}$ G.




FIG. 3: Electronic distribution of the molecular ion $1\sigma_g$ in its ground state in a magnetic field $B = 4.414 \times 10^{13}$ G, with $B_0 = 2.35 \times 10^9$ G, in linear and parallel configuration to the direction of the magnetic field, considering that $R_1 = R_3$ and $R_2 = R_4$ (see Fig. 1).

B. The Excited State $1\pi_u$

The calculations were made for the two values of the conversion factor B_0 and it is shown in Table III. Also for this state, each Ansatz presents a minimum in the energy curve for the finite parameter values $R_{1,\dots,4}$.

For the excited state $1\pi_u$ the electronic distribution is shown in Fig. 4. The positions of the protons are indicated by points in the plane (x, z) . The fact that apparently there are only three points present and not four, it is due to the fact that the distances between the internal protons is very small in comparison with the whole dimension of the system (see Table III). This electronic distribution presents seven maximums (Fig. 4(b)) located in neighboring regions to the protons. From these, those that surround internal protons (see Fig. 1) are $\sim 10^8$ times more pronounced than the minors of them. A comparison of the total energy and the equilibrium distances [14] between molecular ions H_4^{3+} and H_2^+ , both for the excited state $1\pi_u$, shows that both are very similar (see Table IV), indicating that $H_4^{3+} \rightarrow H_2^+ + 2p$ could be a possible channel of decay of the molecular system H_4^{3+} . If a comparison of the total energy of the systems

H_4^{3+} and H_3^{2+} is carry out (Table IV), we found that the molecular ion H_4^{3+} could have a possible channel of decay $H_4^{3+} \rightarrow H_3^{2+} + p$. Among the systems H , H_2^+ , H_3^{2+} and H_4^{3+} , the first of them, H , results as the one with the greatest total energy in the rank of the magnetic fields $0 - 4.414 \times 10^{13}$ G, for which a possible decay $H_4^{3+} \rightarrow H + 3p$ is discarded.

Finally, it is important to mention that if we relax the symmetry conditions ($R_{1,\dots,4}$ are arbitrary) in the minimization, the system returns to the symmetric configuration. This is fulfilled for the ground state ($m = 0$) and for the excited state ($m = -1$).

VII. CONCLUSIONS

Applying the variational method, a detailed study of the $1\sigma_g$ and $1\pi_u$ electronic states of the exotic molecular ion H_4^{3+} in a strong magnetic field was carried out. The magnetic field B , was considered to be constant and homogeneous.

$B(\text{G})$	$B_0(\times 10^9 \text{ G})$	$E_T(\text{Ry})$	$E_b(\text{Ry})$	$R_1(\text{a.u.})$	$R_2(\text{a.u.})$
4.414×10^{13}	2.35	18742.8185	40.1603	0.0651	5.5534
	2.3505	18738.8254	40.1578	0.0651	5.5537

TABLE III: Results for the excited state $1\pi_u$ of the molecular ion H_4^{3+} . For a magnetic field $B = 4.414 \times 10^{13} \text{ G}$, with $B_0 = 2.35 \times 10^9 \text{ G}$ in linear, parallel and symmetric configuration.

	State	$E_T(\text{Ry})$	$E_b(\text{Ry})$	$R_{eq}(\text{a.u.})$
H_2^+	$1\sigma_g$	18728.477	54.5018	0.1016
	$1\sigma_u$	18750.07	32.912	2.021
	$1\pi_u$	18741.89	41.09	0.130
	$1\pi_g$	18757.273	25.7054	2.237
	$1\delta_g$	18747.527	35.407	0.148
	$1\delta_u$	18761.18	21.80	2.230
	$2\sigma_g$	18781.576	1.402	3.120
H_3^{2+}	$1\sigma_g$	18727.7475	55.2312	0.110
	$1\sigma_u$	Not exist		
	$1\pi_u$	18742.7564	40.2223	0.145
	$1\pi_g$	Not exist		
	$1\delta_g$	18748.9067	34.0720	0.167
	$1\delta_u$	Not exist		
H_4^{3+}	$1\sigma_g$	18739.2526	43.7261	0.1149
	$1\pi_u$	18742.8185	40.1603	0.1302

TABLE IV: Comparison between molecular systems with one electron: H_2^+ [8], H_3^{2+} [9] and H_4^{3+} (present) in different states. All of them in linear, symmetric and parallel configuration in the presence of a magnetic field $B = 4.414 \times 10^{13} \text{ G}$ ($B_0 = 2.35 \times 10^9 \text{ G}$). In H_3^{2+} the equilibrium distance R_{eq} corresponds to the distance between to adjacent protons. For the system H_4^{3+} this was considered as the separation between the two internal protons $R_{eq} = 2R_1$ (see Fig. 1).

A. Ground State $1\sigma_g$

An improvement in the variational energy of $\sim 0.2\%$ was obtained in comparison with the first calculations

presents in the literature (Turbiner-López). The equilibrium distances in both case are very similar. This in the symmetric configuration. An important remark is that the system H_4^{3+} can already exist for a magnetic field of intensity $B = 3 \times 10^{13} \text{ G}$.

B. Excited State $1\pi_u$

The existence of the excited state of the molecular ion H_4^{3+} in a magnetic field $B = 4.414 \times 10^{13} \text{ G}$ is seem possible. This state presents a peculiarity in its equilibrium distances (see Table III). R_2 turns out to be ~ 80 times R_1 . The fact that the total energy of the molecular ions H_4^{3+} , H_3^{2+} and H_2^+ is very similar, indicates that the ion H_4^{3+} is very unstable in this state. In this sense it is very unstable since any small disturbance could originate a decay. For which the existence of other excited states does not seem possible, at least for this magnetic field value.

Acknowledgments: The author wants to express his deep gratitude to Dr. A. Turbiner for the introduction to the subject and the constant supervision during the investigations that led to the Bachelor thesis which is based on the results of the present work. I deeply indebted to Dr. J.C. López Vieyra for many valuable discussions and a help in preparation of the manuscript. As well I would like to thank Dr. N. Guevara L. for his help in creating the computer code and his numerous comments during the time when the work was done.

A financial support provided by the CONACYT through the project No. 36650-E is gratefully acknowledged.

-
- [1] Landau. L. *Soviet Z. Soujetunion*, **1** (1991) 185.
 - [2] B. B. Kadomtsev, and V. S. Kudryavtsev. Molecules in an ultrastrong magnetic field. *Pis'ma Zh. Eksp. Teor Fiz. [Sov. Phys. JETP Lett.]*, **13**(1) (1971) 15-19.
 - [3] M. Ruderman. Matter in Superstrong Magnetic Fields: The Surface of a Neutron Star. *Phys. Rev. Lett.* **27** (1971) 1306-1308.
In *IAU Symp. Physics of Dense Matter*, **53** (1974) 117 (ed. by C.J. Hansen, Dordrecht).
 - [4] Turbiner, A. V. and López-Vieyra, J. C. A Hydrogenic Molecular Atmosphere of a Neutron Star. *Mod. Phys. Lett. A*, **19** (2004) 1919-1923.
 - [5] Alexander V. Turbiner and Juan Carlos López Vieyra. One-electron Molecular Systems in a Strong Magnetic Field. *Phys. Reports* **424** (2006) 309-396.
 - [6] J. C. López-Vieyra and A. V. Turbiner. One-electron systems in a strong magnetic field. *Phys. Rev A*, **62** (2000) 022510.
 - [7] H. Olivares Pilón. The Exotic Molecular Ion H_4^{3+} in a Strong Magnetic Field, B.Sc. Thesis, UNAM, Mexico (in Spanish), 2005.
 - [8] A. V. Turbiner and J. C. López-Vieyra. H_2^+ ion in a strong magnetic field. Lowest excited states. *Phys. Rev. A* **69** (2004) 053413.
 - [9] A. V. Turbiner, J. C. López-Vieyra and N. L. Guevara. Exotic H_3^{2+} ion in a strong magnetic field: Linear con-




FIG. 4: (a) Electronic distribution of the molecular ion H_4^{3+} for the excited state $1\pi_u$ in a magnetic field $B = 4.414 \times 10^{13}$ G, with the conversion factor $B_0 = 2.35 \times 10^9$ G. The points indicate the protons position (see text). (b) An enlargement of 10^7 in vertical scale allows the location of the other six pairs of peaks. The two most steep are distinguished here with the numbers 1 and 2, and are indicated with the symbol $\blacklozenge$. For a better visualization, a perpendicular view to the plane (x, z) is presented. Normalization is not fixed.

figuration. *Phys. Rev. A* . **72** (2005) 023403.

- [10] L. D. Landau and E. M. Lifshitz. *Quantum Mechanics* (Pergamon Press Oxford - New York - Toronto - Sydney - Paris - Frankfurt 1977).
- [11] This potential corresponds to the first term of the trial function (Eq. (4a)) with parameters α without permute.
- [12] The binding energy is the necessary energy to separate

the system in its different components.

- [13] Although the first calculations for this system are presented in [6], the comparison is made with the results not published by the same authors.
- [14] This comparison is carry out considering only the distance of the internal protons for the system H_4^{3+} .